

Comentarios

J. M. González Casado.—¿Existen estructuras de cizalla dúctil sin D₂, a los dos lados de la falla frágil?

Respuesta.—Sí existen estructuras de cizalla dúctil sin D₂ en los dos bloques de falla. Los situados en el bloque de techo aparecen en el núcleo de la antiforme de El Cardoso y se observan tanto en los materiales inmediatamente superiores a los gneises como en los inferiores, donde alcanzan un mayor desarrollo.

J. M. González Casado.—¿El modelo de la falla dúctil sin D₂, con movimiento hacia el E y una falla normal posterior, no es el propuesto por González Casado, 1986?

Respuesta.—El modelo propuesto por

González Casado (1986) es, con pequeñas modificaciones, muy similar al propuesto por autores anteriores: Capote *et al.* (1977), González Lodeiro (1981), entre otros. La única diferencia está en que en tu modelo se habla de la existencia de una falla normal, ahora bien, el salto de dicha falla, aunque no está evaluado, no es suficiente para relacionar la zona de cizalla de Berzosa con la que aparece en el núcleo del antiforme de El Cardoso. De hecho, en el modelo propuesto por ti la zona de cizalla de Berzosa se situaría por encima del antiforme de El Cardoso. El modelo aquí propuesto, por el contrario, considera que la llamada zona de cizalla de Berzosa estaría situada a una mayor profundidad de la que actualmente aflora y estaría por debajo del antiforme de El Cardoso y del sinforme de Majael-

rayo, pliegues que, por otra parte, en tu modelo, como en los modelos anteriores, son de F₁, mientras que en el ahora propuesto son de F₃.

C. Casquet.—La falla sería, en tu opinión, frágil y por consiguiente postmetamórfica. Debería de tratarse, pues de una banda cataclástica de relativa importancia en la que dominarían los mecanismos de deformación por ruptura. ¿Dónde se sitúa dicha banda cataclásica al N. de la Cabrera? Toda la deformación que yo he visto hasta ahora es dúctil y sinmetamórfica.

Respuesta.—Sí, y de hecho son muy frecuentes las estructuras cataclásicas que se sobreimponen a estructuras dúctiles de F₂ que, por supuesto, son muy abundantes.

Lacustrine diatomite deposits in the Madrid Basin (Central Spain)

J. P. Calvo. Departamento de Petrología. Facultad de Ciencias Geológicas. Universidad Complutense, 28040 Madrid.

M. Pozo. Departamento de Química Agrícola, Geología y Geoquímica. Facultad de Ciencias. Universidad Autónoma de Madrid. 28049 Madrid.

S. Servant-Vildary. ORSTOM, Laboratoire de Géologie. Muséum National d'Histoire Naturelle, 43, rue de Buffon. 75006 París.

RESUMEN

La parte superior de la Unidad Intermedia del Mioceno en el sector occidental de la Cuenca de Madrid contiene varios niveles diatomíticos, que consisten tanto en diatomitas casi puras como en calizas y margas con contenido variable en diatomeas. Estos niveles fueron depositados en un ambiente lacustre somero caracterizado por condiciones de salinidad moderada y alcalinidad variable, tal como atestiguan la mineralogía de los depósitos y las asociaciones de diatomeas reconocidas. Dentro de éstas se ha distinguido dos asociaciones características, una con predominio de Fragilaria bituminosa y otra constituida por varias especies de Cymbella. Ambas asociaciones se sustituyen entre sí en la vertical de las secciones, indicando fluctuaciones en las condiciones de depósito.

Calvo, J. P.; Pozo, M., y Servant-Vildary, S. (1988): Lacustrine diatomite deposits in the Madrid Basin (Central Spain). *Geogaceta*, 4, 14-17.

Key words: Diatomite, Lacustrine sedimentation, Neogene, Madrid Basin.

Introduction

The occurrence of diatoms within sedimentary deposits of the Madrid Basin (central Spain) has been pointed out only since a few years (Bustillo, 1984). Later, Pozo *et al.* (1985) and Leguey *et al.* (1985) have described true diatomite beds to the North of Esquivias (fig. 1). Here, the diatomites are interbedded within a widely chertified carbonate-clayey succession that has mostly interpreted as lacus-

trine deposits. Most recent studies on these sediments have been devoted to their geochemistry (Bustillo & Bustillo, 1987).

The present work deals with the sedimentological analysis of the diatomite occurrences as well as the preliminary description and paleoecological inferences suited from the study of the diatom assemblages at the Esquivias location. Hitherto, no specific attention has been paid to diatoms themselves in the Neogene record of

the Madrid Basin, probably due either to their scarcity or lacking of specialists to carry out the study. Nevertheless, the analysis of diatoms reveals importance for a comprehensive design of the lacustrine systems during the Neogene.

Stratigraphic location of the diatomite beds

The Miocene formations in the Madrid Basin have been divided into




Fig. 1.—Geographic and geologic setting.

three tecto-sedimentary units (Junco & Calvo, 1983). Diatomite beds occur at the uppermost part of the Intermediate Unit (Middle Aragonian to Lower Vallesian). In the studied area (fig. 1) that unit mainly consists of basal fluvio-lacustrine sediments (micaceous sandstones and mudstones) that laterally interfinger, and are capped by, carbonates and clays of lacustrine origin. The thickness of the unit reaches up near 60 m in this area.

The diatomite beds were recognized in an extensively quarried zone (Malcavado). Two sections were investigated and carefully sampled for further analysis (fig. 2):

— The ESP section is made up by a 1.20 m thick sequence of roughly massive to laminar, powdery white-cream carbonates that overlie a well-defined, tabular chert bed (fig. 2).

— the ESC section consists of a 2.15 m thick sequence of soft, powdery white-cream carbonates that intercalate a thin level of green clays (fig. 2). This section is located a few tens of meters to the North of the ESP one. Although geometric continuity does not exist between the two outcrops, a similar stratigraphic position can be deduced for them by analysing similarities of the underlying sediments.

Mineralogy and petrology

In a previous work, Pozo *et al.* (1985) carried out detailed mineralo-

gical and petrological analysis of the miocene sediments in the Malcovado area. Some of the diatomite beds that we study in this paper (ESP section) were included as level IV within their S-2 section.

A closed sampling in the diatomite beds of the ESC section shows that both their mineralogical and petrological characteristics fluctuate from

level to level (fig. 2), this suggesting sedimentological variations during the sedimentation of the diatomites. Most important changes in mineralogy concern to relative amounts of biogenous opal, calcite, and phyllosilicates (fig. 3). Highest values of opal have been determined in samples 3, 4, 5A and 5B, where highest concentration of diatom frustules can be observed as well.

Petrographically, two main types of diatomite sediments have been recognized: 1) diatomaceous marls and limestones, and 2) clayey to nearly pure diatomites. The former consist of a rather homogeneous mixture of micrite or microsparite, and diatom frustules. The diatom frustules are frequently calcitized and/or partially dissolved. Ostracod carapaces, oogonia and stems of carophytes, and rare tests of gastropods are commonly recognized within the diatomaceous marls and limestones. The second type is seen on thin-section like homogeneous aggregates of diatom frustules showing different sizes, siliceous spicules, and probable phytoliths. Variable amounts of phyllosilicates occur mixed with the siliceous skeletons. The distinction between the two types of diatomaceous sediments clearly arises after




Fig. 2.—Sedimentary logs from the studied sections, with indication of mineralogy and zones defined by diatom assemblages (only for ESC section). Letters for diatom species in ESP section represent: a) *Fragilaria bituminosa*; b) *Cymbella leptoceros*; c) *Navicula halionata*; d) *Anomoeoneis sphaerophora*; e) *Amphora veneta*; f) *Navicula cuspidata*.




Fig. 3.—X-ray diffraction patterns from bulk samples (powder method). A) Diatomaceous limestone. B) Clayey diatomite. C) Diatomite. Symbols: CA: Calcite. Op: Opal. Phy: Phyllosilicates. Fd: Feldspars. Q: Quartz.

their recognition under the microscope or XRD analysis (fig. 3), though it is difficult to stay in outcrop.

Diatoms

Samples collected in the Malcova-deso area were analysed for determination of fossil diatom assemblages. Emphasis was done to the analysis of the ESC section because its greater diversity of flora and more frequent changes throughout the profile, the ESP section being rather homogeneous and dominated by a few species: *Fragilaria bituminosa*, *Cymbella leptoceros*, *Navicula halionata*, *Anomoeoneis sphaerophora*, *Amphora veneta*, and *Navicula cuspidata*. Anyway, a more detailed report on the total diatom content will be offered in a future paper.

The determination of diatoms was realized by standard techniques of sample concentration and further examination under microscope. For each slide, we made a census of diatom species present by counting a hundred specimens. The resulting data show that a rather high variety of diatom species (a total of 50 different species) occurs in the ESC section. However, only a maximum of ten species are present in most of our

samples or occur in significant amounts in some of them (fig. 2). Figure 4 shows some selected specimens of diatoms occurring in the studied deposits.

As within section ESP, diatom flora of section ESC is characterized by the presence of *Fragilaria bituminosa* and its variety *F. bituminosa f. minor*, which are dominant both at the bottom (Zone I) and the top (Zone VI) of the section (fig. 2). In contrast, Zones III, IV, and V show a considerable decrease of *Fragilaria bituminosa*, this species being replaced by *Cymbella leptoceros*, *Navicula radiosa tenella* (Zone III), *Cymbella lapponica*, *Cymbella affinis* (Zone IV), and *Cocconeis placentula*, *Nitzschia epiphytica* (Zone V). Zone II is envisaged as a transition zone, mostly characterized by a high diversity of diatom species.

Fragilaria bituminosa Pantocsek, the most abundant diatom species recognized both in ESC and ESP sections, is a fossil diatom which was found




Fig. 4.—Microphotographs of diatom species. A) *Fragilaria bituminosa* Pantocsek (f. minor). B) *Fragilaria bituminosa* Pantocsek. C) *Amphora veneta* Kützing. D) *Navicula tuscula* (Ehr.) Grunow. E) *Cymbella leptoceros* (Ehr.) Grunow. F) *Navicula brasiliiana* (Cleve) Cleve. Magnification, x1000.

initially in Brackwasser Sarmatian deposits in Hungary (Pantocsek, 1889). Its presence serves to attest the Miocene age of the diatomite deposits in the Madrid Basin.

Paleoenvironment of diatomite sedimentation

The continental character, inferred from regional geology, of the miocene formations, is also attested by diatoms provided a complete absence of non-continental diatom species in the studied samples. The sedimentation of the Intermediate Unit in the Esquivias area began with fluvio-lacustrine deposits that vertically gave pass to a complex mosaic of clays, carbonates, and siliceous sediments that were deposited in lacustrine environments. According to this scheme, diatomites would be deposited during stages of maximum regional enlargement of the lake areas.

Sedimentological data as well as evidence provided by diatoms strongly suggest that, as a whole, deposition took place in shallow lakes and surrounding paludal areas. Probably, the lacustrine sedimentary sequence resulted from small, more or less permanent shallow lakes that shifted, and sometimes coalesced, through time. In this picture, diatomite beds should be deposited into most stable lake areas. Evidence of very shallow conditions, such as root holes, suggest that the lake experienced some oscillations of the water level. Further evidence of shallowness is provided

by diatoms. The whole of the diatom specimens that we have recognized in our samples is constituted by epiphytic and/or epilithic species. Living forms of these species are commonly found in shallow waters, frequently in littoral lake environments.

A closer view of the distribution of diatoms in vertical sequence (Zones I to VI in section ESC) allows to infer that Zones I and VI, both of them characterized by *Fragilaria bituminosa*, reflect slightly higher salinities in lake water. This circumstance is clearer in Zone VI, where *Fragilaria bituminosa* occurs associated with mesohalobien species, such as *Anomoeoneis sphaerophora*, *Navicula cincta*, *Amphora veneta*, and *Amphora coffaeiformis*. On the other hand, Zones III, IV, and V are characterized by *Cymbella* species, which are oligohalobien and alkaliphilic species, and they are indicative of fresh-water, shallow lake environments. In this context, Zone V displaying the highest occurrence of *Coccconeis placentula*, represents a more marked lowstand in water level. Finally, the transitional Zone II would represent an episode of progressive freshening between Zones I and III.

It follows from the previous paragraph that diatoms in the miocene formations of the Malcovadeso area are mostly freshwater or tolerant of moderately saline conditions. This seems to fit well with the commonly observed low-Mg calcite composition of the carbonate phase in the diatomite beds. In addition, diatom ecology indicates that the observed species were adapted to somewhat alkali-

ne waters, which in turn favoured the formation of authigenic clays such as palygorskite and sepiolite (Leguey et al., 1985) within the shallow lake. Authigenesis of these clays could be related to the colloidal silica provided from dissolution of diatom frustules.

Finally, main diagenetic features in the diatomites concern to corrosion or even total dissolution and later calcitization of the diatom frustules. Nevertheless, the definite paths of these processes have not been yet evaluated. It does not seem unreal to assume that diatom dissolution mostly accounted for the formation of the opaline levels widely extended within the miocene formations in the studied area.

References

- Bustillo, M. A. (1984): *Estudios geol.*, 40, 137-151.
- Bustillo, M. A. y Bustillo, M. (1987): *Resúmenes II Congreso Nacional de Geoquímica*, Soria, 183-185.
- Junco, F. y Calvo, J. P. (1983): en, *Geología de España, Libro Jubilar J. M. Ríos*, t. II, 534-543.
- Leguey, S.; Pozo, M. y Medina, J. A. (1985): *Miner. Petrogr. Acta*, 29-A, 287-301.
- Pantocsek, J. (1889): *Beiträge zur Kenntnis der Fossilien Bacillarien Hongarns*. Vol. II, *Brackwasser Bacillarien*, 123 pp. Nagy Tapolcsany, Julius Platzko.
- Pozo, M.; Medina, J. A. y Leguey, S. (1985): *Bol. Soc. Esp. Mineralogía*, 271-183.

Recibido el 8 de enero de 1988
Aceptado el 21 de enero de 1988

Isótopos estables de carbono y oxígeno de los rellenos de calcita (falsa ágata) y su interés paleoambiental

C. Jiménez de Cisneros. E. E. Zaidín. CSIC. Albareda, 1. Granada.

J. Linares. E. E. Zaidín. CSIC. Albareda, 1. Granada.

E. Reyes. E. E. Zaidín. CSIC. Albareda, 1. Granada.

J. A. Vera. Departamento Estratigrafía-Paleontología. Universidad de Granada.

ABSTRACT

The isotopic composition of modern calcite fills (onyx marble) in fractures and related caves is studied. Carbon and oxygen stable isotopes in this material are similar to the vadose speleothems (stalactites and stalagmites) in modern caves and to calcareous concre-